

List of Supplementary Figures of

Transport Patterns of Global Aviation NO_x and their Short-term O₃ Radiative Forcing – A Machine Learning Approach

Jin Maruhashi et al.

Correspondence to: Irene C. Dedoussi (I.C.Dedoussi@tudelft.nl)

Figure	Description
S1	Temporal evolution of the air parcel trajectory pressure altitude of median trajectories (dots in red) based on 50 Lagrangian paths (dots in blue) for all 28 emission points for North America for January – March 2014.
S2	Horizontal trajectories for air parcels released in North America during January 2014. The black dots denote the emission starting points. The red dots represent the median trajectory while the blue dots represent the 50 Lagrangian trajectories for each of the 28 emission points.
S3	Determination of optimal cluster threshold θ for North America during January – March 2014. Vertical axis is the sum of the averaged distance $E(\theta)$ as a function of the number of clusters (blue curve). Based on the L-method, the optimal cluster number is found via the intersection of two linear regressions (red).
S4	Vertical profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during January 2014 for all emission points in North America.
S5	Horizontal profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during January 2014 for all emission points in North America.
S6	Horizontal profiles of mean RF in $mW \cdot m^{-2}$ during January 2014 for all 28 emission points in North America.
S7	Temporal evolution of the air parcel trajectory pressure altitude of median trajectories (dots in red) based on 50 Lagrangian paths (dots in blue) for all 28 emission points for North America for July – September 2014.
S8	Horizontal trajectories for air parcels released in North America during July 2014. The black dots denote the emission starting points. The red dots represent the median trajectory while the blue dots represent the 50 Lagrangian trajectories for each of the 28 emission points.
S9	Determination of optimal cluster threshold θ for North America during July – September 2014. Vertical axis is the sum of the averaged distance $E(\theta)$ as a function of the number of clusters (blue curve). Based on the L-method, the optimal cluster number is found via the intersection of two linear regressions (red).
S10	Vertical profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during July 2014 for all emission points in North America.
S11	Horizontal profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during July 2014 for all emission points in North America.
S12	Horizontal profiles of mean RF in $mW \cdot m^{-2}$ during July 2014 for all 28 emission points in North America.
S13	Temporal evolution of the air parcel trajectory pressure altitude of median trajectories (dots in red) based on 50 Lagrangian paths (dots in blue) for all 28 emission points for South America for January – March 2014.
S14	Horizontal trajectories for air parcels released in South America during January 2014. The black dots denote the emission starting points. The red dots represent the median trajectory while the blue dots represent the 50 Lagrangian trajectories for each of the 28 emission points.
S15	Determination of optimal cluster threshold θ for South America during January – March 2014. Vertical axis is the sum of the averaged distance $E(\theta)$ as a function of the number of clusters (blue curve). Based on the L-method, the optimal cluster number is found via the intersection of two linear regressions (red).
S16	Vertical profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during January 2014 for all emission points in South America.
S17	Horizontal profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during January 2014 for all emission points in South America.
S18	Horizontal profiles of mean RF in $mW \cdot m^{-2}$ during January 2014 for all 28 emission points in South America.
S19	Temporal evolution of the air parcel trajectory pressure altitude of median trajectories (dots in red) based on 50 Lagrangian paths (dots in blue) for all 28 emission points for South America for July – September 2014.
S20	Horizontal trajectories for air parcels released in South America during July 2014. The black dots denote the emission starting points. The red dots represent the median trajectory while the blue dots represent the 50 Lagrangian trajectories for each of the 28 emission points.

S21	Determination of optimal cluster threshold θ for South America during July – September 2014. Vertical axis is the sum of the averaged distance $E(\theta)$ as a function of the number of clusters (blue curve). Based on the L-method, the optimal cluster number is found via the intersection of two linear regressions (red).
S22	Vertical profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during July 2014 for all emission points in South America.
S23	Horizontal profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during July 2014 for all emission points in South America.
S24	Horizontal profiles of mean RF in $mW \cdot m^{-2}$ during July 2014 for all 28 emission points in South America.
S25	Temporal evolution of the air parcel trajectory pressure altitude of median trajectories (dots in red) based on 50 Lagrangian paths (dots in blue) for all 28 emission points for Eurasia for July – September 2014.
S26	Horizontal trajectories for air parcels released in Eurasia during July 2014. The black dots denote the emission starting points. The red dots represent the median trajectory while the blue dots represent the 50 Lagrangian trajectories for each of the 28 emission points.
S27	Determination of optimal cluster threshold θ for Eurasia during July – September 2014. Vertical axis is the sum of the averaged distance $E(\theta)$ as a function of the number of clusters (blue curve). Based on the L-method, the optimal cluster number is found via the intersection of two linear regressions (red).
S28	Vertical profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during July 2014 for all emission points in Eurasia.
S29	Horizontal profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during July 2014 for all emission points in Eurasia.
S30	Horizontal profiles of mean RF in $mW \cdot m^{-2}$ during July 2014 for all 28 emission points in Eurasia.
S31	Temporal evolution of the air parcel trajectory pressure altitude of median trajectories (dots in red) based on 50 Lagrangian paths (dots in blue) for all 28 emission points for Eurasia for January – March 2014.
S32	Horizontal trajectories for air parcels released in Eurasia during January 2014. The black dots denote the emission starting points. The red dots represent the median trajectory while the blue dots represent the 50 Lagrangian trajectories for each of the 28 emission points.
S33	Determination of optimal cluster threshold θ for Eurasia during January – March 2014. Vertical axis is the sum of the averaged distance $E(\theta)$ as a function of the number of clusters (blue curve). Based on the L-method, the optimal cluster number is found via the intersection of two linear regressions (red).
S34	Vertical profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during January 2014 for all emission points in Eurasia.
S35	Horizontal profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during January 2014 for all emission points in Eurasia.
S36	Horizontal profiles of mean RF in $mW \cdot m^{-2}$ during January 2014 for all 28 emission points in Eurasia.
S37	Temporal evolution of the air parcel trajectory pressure altitude of median trajectories (dots in red) based on 50 Lagrangian paths (dots in blue) for all 28 emission points for Africa for January – March 2014.
S38	Horizontal trajectories for air parcels released in Africa during January 2014. The black dots denote the emission starting points. The red dots represent the median trajectory while the blue dots represent the 50 Lagrangian trajectories for each of the 28 emission points.
S39	Determination of optimal cluster threshold θ for Africa during January – March 2014. Vertical axis is the sum of the averaged distance $E(\theta)$ as a function of the number of clusters (blue curve). Based on the L-method, the optimal cluster number is found via the intersection of two linear regressions (red).
S40	Vertical profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during January 2014 for all emission points in Africa.
S41	Horizontal profiles of mean O_3 disturbances in $nmol \cdot mol^{-1}$ during January 2014 for all emission points in Africa.

S42	Horizontal profiles of mean RF in $\text{mW}\cdot\text{m}^{-2}$ during January 2014 for all 28 emission points in Africa.
S43	Temporal evolution of the air parcel trajectory pressure altitude of median trajectories (dots in red) based on 50 Lagrangian paths (dots in blue) for all 28 emission points for Africa for July – September 2014.
S44	Horizontal trajectories for air parcels released in Africa during July 2014. The black dots denote the emission starting points. The red dots represent the median trajectory while the blue dots represent the 50 Lagrangian trajectories for each of the 28 emission points.
S45	Determination of optimal cluster threshold θ for Africa during July – September 2014. Vertical axis is the sum of the averaged distance $E(\theta)$ as a function of the number of clusters (blue curve). Based on the L-method, the optimal cluster number is found via the intersection of two linear regressions (red).
S46	Vertical profiles of mean O_3 disturbances in $\text{nmol}\cdot\text{mol}^{-1}$ during July 2014 for all emission points in Africa.
S47	Horizontal profiles of mean O_3 disturbances in $\text{nmol}\cdot\text{mol}^{-1}$ during July 2014 for all emission points in Africa.
S48	Horizontal profiles of mean RF in $\text{mW}\cdot\text{m}^{-2}$ during July 2014 for all 28 emission points in Africa.
S49	Temporal evolution of the air parcel trajectory pressure altitude of median trajectories (dots in red) based on 50 Lagrangian paths (dots in blue) for all 28 emission points for Australasia for July – September 2014.
S50	Horizontal trajectories for air parcels released in Australasia during July 2014. The black dots denote the emission starting points. The red dots represent the median trajectory while the blue dots represent the 50 Lagrangian trajectories for each of the 28 emission points.
S51	Determination of optimal cluster threshold θ for Australasia during July – September 2014. Vertical axis is the sum of the averaged distance $E(\theta)$ as a function of the number of clusters (blue curve). Based on the L-method, the optimal cluster number is found via the intersection of two linear regressions (red).
S52	Vertical profiles of mean O_3 disturbances in $\text{nmol}\cdot\text{mol}^{-1}$ during July 2014 for all emission points in Australasia.
S53	Horizontal profiles of mean O_3 disturbances in $\text{nmol}\cdot\text{mol}^{-1}$ during July 2014 for all emission points in Australasia.
S54	Horizontal profiles of mean RF in $\text{mW}\cdot\text{m}^{-2}$ during July 2014 for all 28 emission points in Australasia.
S55	Temporal evolution of the air parcel trajectory pressure altitude of median trajectories (dots in red) based on 50 Lagrangian paths (dots in blue) for all 28 emission points for Australasia for January – March 2014.
S56	Horizontal trajectories for air parcels released in Australasia during January 2014. The black dots denote the emission starting points. The red dots represent the median trajectory while the blue dots represent the 50 Lagrangian trajectories for each of the 28 emission points.
S57	Determination of optimal cluster threshold θ for Australasia during January – March 2014. Vertical axis is the sum of the averaged distance $E(\theta)$ as a function of the number of clusters (blue curve). Based on the L-method, the optimal cluster number is found via the intersection of two linear regressions (red).
S58	Vertical profiles of mean O_3 disturbances in $\text{nmol}\cdot\text{mol}^{-1}$ during January 2014 for all emission points in Australasia.
S59	Horizontal profiles of mean O_3 disturbances in $\text{nmol}\cdot\text{mol}^{-1}$ during January 2014 for all emission points in Australasia.
S60	Horizontal profiles of mean RF in $\text{mW}\cdot\text{m}^{-2}$ during January 2014 for all 28 emission points in Australasia.