


Chapter 9

Supplemental Data 4: Raw figures pituitary Cushing only


Generic quality of life in untreated Cushing disease patients.



Higher score represents lower quality of life: NHP, SCL-90-R

Partially overlapping populations: Milian 2013 and Milian 2013 (2)


Generic quality of life in treated Cushing disease patients.



Higher score represents lower quality of life: GHQ-28, NHP, PSI (except well-being), SCL-90(-R), SF-20, SIP


Partially overlapping populations: Milian 2013, Milian 2013 (2), and Milian 2014; Milian 2015 and Psaras 2011

Generic quality of life in Cushing disease patients before versus after treatment.




Partially overlapping populations: Milian 2013 and Milian 2013 (2)

Disease specific quality of life in untreated Cushing disease patients.




Higher score represents lower quality of life: Tuebingen CD-25

Disease specific quality of life in treated Cushing disease patients.




Higher score represents lower quality of life: Tuebingen CD-25

Disease specific quality of life in Cushing disease patients before versus after treatment.




Quality of life: domain anxiety, in treated Cushing disease patients.




Higher score represents lower quality of life: BAI, FQ, HADS, MASQ-30, STAI
 Partially overlapping populations: Andela 2013, Tiemensma 2010, and Van Aken 2005

Quality of life: domain anxiety, in Cushing disease patients after treatment versus healthy controls.



Partially overlapping populations: Andela 2013, Tiemensma 2010, and Van Aken 2005


Quality of life: domain depression, in treated Cushing disease patients.



Higher score represents lower quality of life: BDI(-II), HADS, HRSD, IDS, MADRS


Partially overlapping populations: Andela 2013, Tiemensma 2010, and Van Aken 2005

Quality of life: domain depression, in Cushing disease patients after treatment versus healthy controls.




Partially overlapping populations: Andela 2013, Tiemensma 2010, and Van Aken 2005

Quality of life: domain fatigue, in treated Cushing disease patients.




Higher score represents lower quality of life: CIS, MFI-20

Cognitive function: executive functioning, in untreated Cushing disease patients.




Higher score represents worse cognitive function: TMT

Cognitive function: executive functioning, in treated Cushing disease patients.




Higher score represents worse cognitive function: TMT


Cognitive function: executive functioning, in Cushing disease patients before versus after treatment.




Cognitive function: executive functioning, in Cushing disease patients after treatment versus healthy controls.




Cognitive function: attention, in treated Cushing disease patients.




Cognitive function: attention, in Cushing disease patients after treatment versus healthy controls.




Quality of life: SF-36, in untreated Cushing disease patients.




Quality of life: SF-36, in treated Cushing disease patients.




Quality of life: SF-36, in Cushing disease patients before versus after treatment.



Quality of life: SF-36, in Cushing disease patients after treatment versus healthy controls.




Quality of life: SCL-90-R, in untreated Cushing disease patients.




Higher score represents lower quality of life: SCL-90-R

Quality of life: SCL-90-R, in treated Cushing disease patients.




Higher score represents lower quality of life: SCL-90-R

Quality of life: SCL-90-R, in Cushing disease patients before versus after treatment.




Quality of life: GHQ-28, in treated Cushing disease patients.




Higher score represents lower quality of life: GHQ-28


Quality of life: WHOQoL-BREF, in treated Cushing disease patients.



Quality of life: CushingQoL, in treated Cushing disease patients.



Quality of life: HADS, in treated Cushing disease patients.



Higher score represents lower quality of life: HADS